

5      **Costimulating polypeptide of T cells, monoclonal anti-  
bodies, and the preparation and use thereof**

10            The invention relates to a polypeptide (8F4  
molecule) having the biological activity of costimula-  
ting T cells. The invention further relates to monoclo-  
nal antibodies against the 8F4 molecule and hybridoma  
cells which produce the monoclonal antibodies. The in-  
vention additionally relates to the use of substances  
which inhibit the biological activity of the polypepti-  
15      de 8F4 according to the invention, in particular mono-  
clonal antibodies, natural or synthetic ligands, ago-  
nists or antagonists, as pharmaceuticals. In particu-  
lar, the invention relates to the use of these substan-  
ces for the prevention or therapy of disorders in which  
20      the immune system is involved, in particular for the  
treatment of autoimmune diseases and for the prevention  
of rejection reactions with organ transplants. The in-  
vention additionally relates to the use of the 8F4 mo-  
lecle or of cells which contain the 8F4 molecule as  
25      pharmaceuticals, in particular for the prevention or  
therapy of disorders in which the immune system is in-  
volved, in particular for the treatment of cancers,  
Aids, asthmatic disorders or chronic viral diseases  
such as HCV or HBV infections. The invention further  
30      relates to the use of substances which specifically re-  
cognize the polypeptide according to the invention, in  
particular monoclonal antibodies, natural or synthetic  
ligands, agonists or antagonists, for the diagnosis of  
disorders in which the immune system is involved. In  
35      particular, the invention relates to diagnosis by means  
of an ELISA detection, a flow cytometry or a Western  
blot, a radioimmunological detection, a nephelometry or  
a histochemical staining.

40            T lymphocytes recognize their antigen, which  
is presented by "antigen-presenting cells", for example  
dendritic cells, B cells and macrophages, through their  
T-cell receptor. Recognition of the antigen by the  
T-cell receptor alone is, however, in most cases insuf-  
ficient for adequate activation of T lymphocytes. The  
45      latter makes additional simultaneous stimulation (also  
called "costimulation" hereinafter) by other receptor  
molecules on the surface of the T lymphocytes necessa-

ry. One of these receptor molecules is the so-called CD28 receptor which is stimulated by the costimulating molecule B7. If these "costimulatory" molecules, for example CD28, are effective, then the activation of the T cell reaches an adequate level after recognition of the antigen by the T-cell receptor. After such a complete activation, the T cell expresses additional molecules, for example CD25, CD69, CD71, on the surface and synthesizes numerous cytokines, for example IL-2 and IFN- $\gamma$ , which function as messengers. Both these additional surface molecules and the cytokines serve for the T cell to exchange information with other cells in the immune system. The activated T cells direct the entire antigen-specific immune defences through the additional surface molecules and the cytokines. Both the generation of cytotoxic cells ("killer cells") and the generation of antigen-specific antibodies by B cells is controlled in this way. Cytotoxic cells, as well as the specifically formed antibodies, eliminate viral or bacterial pathogens which enter the body. In some cases, however, the immune response goes too far, and the immune system is directed against the body's own cells. This leads to the occurrence of "autoimmune diseases", for example to rheumatoid arthritis, ankylosing spondylitis, Sjögren's syndrome, ulcerative colitis inter alia. One of the essential sites of cooperation between antigen-activated T cells and other cells of the immune system are the secondary lymphatic organs, including the tonsils. This is where the T lymphocytes are activated by the antigen presented by dendritic cells, and this is where T lymphocytes interact with B cells. Through this interaction, B cells secrete, after several intermediate stages of differentiation, antigen-specific antibodies of the IgM and IgG types.

The costimulatory molecule which has been characterized best and is among the most effective to date is the CD28 surface molecule (called CD28 receptor or CD28 hereinafter) which is constitutively expressed on a large fraction of T cells. Costimulation by CD28 *in vitro* leads, after recognition of the antigen by the T-cell receptor, to a very large increase in cytokine secretion, for example of IL-2 and IFN- $\gamma$ , and to a marked up-regulation of the expression of cell surface molecules such as CD25, CD69, CD71, which are necessary for interaction of T cells with other immune cells, for example B lymphocytes; cf. Chambers and Allison, *Current Opinion in Immunology* 9 (1997), 396-404. Costimulation via the CD28 receptor can also markedly increase the proliferation of T lymphocytes. In addition, costi-

mulation via the CD28 receptor optimizes the T-cell control of B-lymphocyte function so that there is increased secretion of antibodies.

If the function of the CD28 receptor is abolished, there is a drastic loss of function in the immune defences. This has been shown by means of a transgenic mouse in which the CD28 gene was destroyed by homologous recombination (a so-called "CD28 knock-out"). The destruction in this way of activation of the antigen-specific T cells leads to lack of costimulation. This in turn leads to a disturbance of T-cell function, that is to say to a reduced proliferation of T cells and to a drastically reduced synthesis of various cytokines. The lack of costimulation eventually leads to a reduced function of the antigen-specific immune defences. Thus, inter alia, the formation of antigen-specific IgG1 and IgG2 antibodies by B lymphocytes is reduced to 10% of the normal level through the lack of CD28; cf. Shahinian et al., *Science* 262 (1993), 609-612; Lucas et al. *Journal of Immunology* 154 (1995), 5757-5768. It is also possible *in vitro* to prevent the Aids virus entering T lymphocytes by costimulation by CD28; cf. Riley et al., *Journal of Immunology* 158 (1997), 5545-5553. Corresponding experiments *in vivo* have not yet been carried out. It is known that CD28 switches on many cytokine genes which may lead to considerable side effects *in vivo*. Blockade of CD28 receptors by a soluble CTLA-4 immunoglobulin molecule has been employed successfully in a monkey model to prevent the rejection of transplanted kidneys. In this case, CTLA-4 had been employed in combination with an antibody against the CD40 ligand molecule; cf. Kirk et al., *Proc. Natl. Acad. Sci. USA* 94 (1997) 8789-8794. However, blockade of CD28 receptors affects all T lymphocytes and not just those already activated because CD28 is constitutively expressed on T lymphocytes.

There is thus a need for a costimulating surface molecule which is expressed only on activated T lymphocytes. The invention is therefore based on the object of providing a surface molecule on activated T cells which has a strong costimulatory effect on central functions of T lymphocytes. Another object of the invention is to provide substances, for example monoclonal antibodies against the costimulatory surface molecule, natural or synthetic ligands, agonists or antagonists of the surface molecule.

In a first embodiment, the invention relates to a polypeptide having the biological activity of co-

stimulation of T cells, characterized in that a) the polypeptide occurs on activated CD4<sup>+</sup> and CD8<sup>+</sup> T lymphocytes but not on resting or activated B cells, granulocytes, monocytes, NK cells (natural killer cells) or dendritic cells, and b) the polypeptide is a dimer, the polypeptide having a molecular weight of about 55 to 60 kDa (kilodalton) determined in a non-reducing sodium dodecyl sulphate polyacrylamide gel electrophoresis (SDS-PAGE), and the two polypeptide chains of the polypeptide having a molecular weight of about 27 kDa and about 29 kDa measured in a reducing SDS-PAGE.

The polypeptide according to the invention (also called 8F4 molecule or 8F4 hereinafter) is expressed only after activation of the T lymphocytes, specifically both on CD4<sup>+</sup> and on CD8<sup>+</sup> T cells. In a non-reducing SDS-PAGE, the 8F4 molecule has a molecular weight between about 55 and 60 kDa (kilodalton). The 8F4 molecule is composed of two peptide chains, and the two peptide chains have a molecular weight of about 27 and about 29 kDa in a reducing SDS-PAGE. The 8F4 antigen can be unambiguously detected histologically on activated T lymphocytes in the lymphatic tissue of the tonsils and lymph nodes, especially in the germinal centres, the site of interaction of T lymphocytes and B lymphocytes in the generation of antibodies. Tonsillar T cells isolated *ex vivo* are about 50-80% positive for the 8F4 antigen and show signs of advanced activation. The 8F4 molecule is not detectable on resting or activated B cells, granulocytes, monocytes, NK cells and dendritic cells.

An important biological activity of the 8F4 molecule is its costimulating activity on T lymphocytes. The costimulating activity can be determined by the method of Linsley et al., *Journal of Experimental Medicine* 176 (1992), 1595-604. The costimulating activity of the 8F4 molecule resembles the costimulating activity of the CD28 molecule, which has been identified as the central enhancement element of antigen recognition by the immune system. The 8F4 molecule differs in many aspects from CD28, however. Thus, expression of the 8F4 molecule on the surface of the T cells requires induction, whereas CD28 is constitutively expressed. There are also distinct differences detectable in the function: costimulation by CD28 leads to overexpression of numerous lymphokines, *inter alia* of interleukin-2 (IL-2). Costimulation by 8F4 also leads to enhanced secretion of lymphokines, but not of IL-2. The costimulatory activity of the 8F4 molecule thus differs from the activity of the CD28 molecule.

Since stimulation by 8F4 does not switch on all cytokine genes, costimulation by 8F4 *in vivo* is advantageous, for example compared with costimulation via the CD28 receptor. Moreover, the induction, the expression, the  
5 site of expression and the function of the 8F4 molecule differ from all other known molecules with costimulatory activity.

The 8F4 molecule according to the invention  
10 is a novel surface molecule on activated T cells which has a strong costimulatory effect on central functions of T lymphocytes. Expression *in vivo* indicates inter alia an essential involvement of the 8F4 molecule in the cooperation of T cells with other cells of the immune system such as B cells or dendritic cells within  
15 the humoral and cellular immune defences against viruses and bacteria.

After expression, the 8F4 molecule has *in vitro* a strong costimulatory effect on various functions of T lymphocytes:

- 20 1. Marked enhancement of the proliferation of T lymphocytes.
2. Marked enhancement of the synthesis of certain cytokines by T lymphocytes.
- 25 3. Greatly increased expression of control molecules, for example surface molecules and cytokines, on and in T lymphocytes.
4. Marked improvement in T-cell-induced antibody formation (IgM and IgG) by B cells.

The present invention furthermore provides a polypeptide having the biological activity of costimulation of T cells and having an amino acid sequence which shows at least 40% homology with the sequence comprising  
30 199 amino acids in Fig. 15 (SEQ ID NO:2), or a biologically active fragment or an analogue thereof. A biologically active fragment or analogue is a fragment or analogue which likewise shows a costimulatory effect on  
35 T-cell lymphocytes or at least displays a biological effect of the nature of a blockage. Preference is given to a polypeptide or a biologically active fragment or analogue thereof which shows at least 60% homology with  
40 the sequence comprising 199 amino acids in Fig. 15 (SEQ ID NO:2). In a particularly preferred embodiment, the polypeptide according to the invention comprises an amino acid sequence which shows at least 80% homology  
45 with the sequence comprising 199 amino acids in Fig. 15

(SEQ ID NO:2), or a biologically active fragment or analogue thereof.

5 A particularly preferred polypeptide has the biological activity of costimulation of T cells and comprises an amino acid sequence as shown in Fig. 15 (SEQ ID NO:2), or a biologically active fragment or an analogue thereof.

10 The invention includes allelic variants, fragments and analogues of the 8F4 molecule. These variants include naturally occurring allelic variants, substitution analogues in which one or more amino acids have been substituted by different amino acids, substitution analogues in which one or more amino acids have been substituted by different amino acids, deletion  
15 analogues in which one or more amino acids have been deleted and addition analogues in which one or more amino acids have been added. Deletion and addition of one or more amino acids may be done either at an internal region of the polypeptide or at the amino or carboxyl terminus.  
20

Polypeptides according to the invention fused to heterologous polypeptides are likewise embraced.

25 In another embodiment, the invention relates to DNA sequences which encode a polypeptide according to the invention or a biologically active fragment or analogue thereof.

30 These DNA sequences include the sequence shown in SEQ ID NO:1 (Fig. 16) as well as allelic variants, fragments, and analogues having biological activity.

A preferred DNA sequence encodes a polypeptide having the biological activity of costimulation of T cells, the sequence being selected from the group consisting of:

- 35 a) the DNA sequence shown in SEQ ID NO:1 (Fig. 16) and its complementary strand  
b) DNA sequence hybridizing with the sequences in (a) and  
40 c) DNA sequences which, because of the degeneracy of the genetic code, hybridize with the sequences in (a) and (b). The aforementioned DNA sequences preferably hybridize together under stringent conditions.

45 Also provided are vectors which comprise these DNA sequences, and host cells which are transformed or transfected with these vectors.

In another embodiment, the invention relates to monoclonal antibodies against the 8F4 molecule. The monoclonal antibodies according to the invention can be prepared in a conventional way by the method described by Milstein and Köhler, *Nature* 256 (1975), 495-497. In particular, the monoclonal antibodies according to the invention can be prepared by immunizing mice with T cells which have been activated *in vitro* with phorbol myristate acetate (PMA) and ionomycin ("2-signal system") for 24 h. The spleen cells of the immunized mice are fused with myeloma cells. 8F4-specific monoclonal antibodies are identified by their recognition of 2-signal-activated but not resting T lymphocytes. Moreover 8F4-specific antibodies do not stain T cells stimulated with one signal (either PMA or ionomycin) in a detection method carried out in a conventional way. 8F4-specific antibodies produce a typical staining pattern of tonsillar T cells and recognize an antigen of about 55 to 60 kDa in a non-reducing SDS-PAGE and of about 27 kDa and about 29 kDa in a reducing SDS-PAGE on activated T lymphocytes.

In another embodiment, the invention relates to hybridoma cells which produce the monoclonal antibodies according to the invention.

In another embodiment, the invention relates to the use of substances which inhibit the biological activity of the polypeptide 8F4 according to the invention as pharmaceuticals. The use of the monoclonal antibodies according to the invention, natural or synthetic ligands, agonists or antagonists of the 8F4 molecule is particularly preferred. These substances can be used as pharmaceuticals for the prevention or therapy of disorders in which the immune system is involved, in particular for the treatment or autoimmune diseases or for prevention of rejection reactions in organ transplants. Blockade of the interaction of the 8F4 antigen with its receptor improves, for example, the prevention of organ rejection because such a blockade affects only previously activated T lymphocytes. Another embodiment of the invention relates to the use of the polypeptide according to the invention as pharmaceutical. The polypeptide according to the invention can be used in particular for the prevention or therapy of disorders in which the immune system is involved, in particular for the treatment of cancers, AIDS, asthmatic disorders or chronic viral diseases such as HCV or HBV infections.

The polypeptide according to the invention can likewise be introduced into cells in a conventional

way so that these cells for example constitutively express the polypeptide. For example, the nucleic acid sequence encoding the polypeptide or a vector comprising the nucleic acid sequencing encoding the polypeptide, for example the cDNA or genomic DNA, promoters, enhancers and other elements required for expression of the nucleic acid sequence can be inserted into a cell. The 8F4 cDNA (2641 nucleotides) depicted in Fig. 16 (SEQ ID NO:1) or fragments or derivatives thereof, is preferably employed for expression of the polypeptide according to the invention or fragments thereof.

The polypeptide according to the invention can also be introduced for example by means of liposomes into cells which then form the polypeptide on their cell surface. These cells can be used as pharmaceuticals according to the invention, in particular for restoring correct regulation of the human immune system, as occurs within the framework of numerous chronic infectious diseases, for example within the framework of AIDS, asthmatic disorders or in chronic viral hepatitis (for example HCV, HBV), or for stimulating the immune system in vitro or in vivo such as, for example, be used for the therapy of cancers.

In another embodiment, substances which specifically recognize the polypeptide according to the invention are used for diagnosing disorders in which the immune system is involved, the substances embracing in particular a monoclonal antibody, natural or synthetic ligands, agonists or antagonists. It is possible to use for the diagnosis for example an ELISA detection, flow cytometry, Western blot, radioimmunoassay, nephelometry or a histochemical staining. The substances which recognize the polypeptide according to the invention also comprise nucleic acid sequences, the latter preferably being employed for hybridization and/or nucleic acid (RNA, DNA) amplification (for example PCR).

In another embodiment, the invention relates to substances which have a positive or negative effect on (modulate) the signal transduction pathway of the polypeptide according to the invention into the T cell, and to the use of these substances as pharmaceuticals.

In another embodiment, the invention relates to substances which prevent up-regulation of the polypeptide according to the invention on the T-cell surface, and to the use thereof as pharmaceuticals.



In another embodiment, the polypeptide according to the invention or fragments thereof is expressed by a transgenic animal.

5 In another embodiment, the invention embraces a transgenic animal in which the gene which codes for the polypeptide according to the invention has been switched off ("knock-out").

The figures serve to illustrate the invention:

10 Fig. 1 shows the result of an immunoprecipitation of the 8F4 antigen from activated human T cells. (a) Sodium dodecyl sulphate polyacrylamide gel electrophoresis (SDS-PAGE; 12% polyacrylamide gel (PAA gel)) reducing, (b) SDS-PAGE (10% PAA gel) non-reducing. The  
15 conditions for elution of the antigen from the 8F4 matrix are indicated. "SDS" means sodium dodecyl sulphate; "DTT" means dithiothreitol, "Mr" means molecular weight and "kDa" means kilodalton.

20 Fig. 2a shows the result of a flow cytometry after induction of the 8F4 antigen on CD4<sup>+</sup> T cells. The activation time for the T cells is indicated in parentheses. "PMA" means phorbol myristate acetate; "PHA" means phytohaemagglutinin; "OKT3" is a monoclonal antibody against CD3; "MLR" means mixed lymphocyte reaction;  
25 "mAK 9.3" is a monoclonal antibody against CD28; "SEB" means staphylococcal enterotoxin B.

Fig. 2b shows the result for the kinetics of induction of the 8F4 antigen on CD4<sup>+</sup> T cells after activation with PMA and ionomycin in a flow cytometry. The  
30 immunofluorescence (log) is plotted against the cell count.

Fig. 3 shows the result of a flow cytometry for identifying molecules which are involved in the induction of 8F4 in the mixed lymphocyte reaction. "bio"  
35 means biotinylated antibody.

Fig. 4 shows the result of a histochemical investigation for localization of 8F4-positive cells in the tonsil.

40 Fig. 5 shows the result of an expression analysis of 8F4 on T and B cells from human tonsils in a flow cytometry. "bioPE" means biotinylated antibody and streptavidin-phycoerythrin secondary reagent.

45 Fig. 6 shows the coexpression of the 8F4 molecule with other activation markers (CD69, CD45) in a flow cytometry.

Fig. 7 shows diagrammatically the enhanced expression of activation molecules on T lymphocytes after costimulation by 8F4. Open circles (O) represent 8F4 antibodies; triangles (♦) represent nonspecific antibodies of the same isotype; filled circles (●) represent anti-CD28 antibodies-9.3.

Fig. 8 shows a diagrammatic comparison of the costimulating effect of 8F4 with the costimulating effect of CD28. "mAk" means monoclonal antibodies; "ATAC" means "activation induced T-cell-derived and chemokine-related"; "cpm" means radioactive disintegrations per minute.

Fig. 9 shows diagrammatically the enhancement of the synthesis of the antibodies of the IgM and IgG types by B cells after costimulation of T cells. "ng" means nanogram; "ml" means millilitre; "mAk" means monoclonal antibody.

Fig. 10 shows diagrammatically the prevention of the activation-induced apoptosis of peripheral T cells after costimulation by 8F4.

Fig. 11 shows expression of the 8F4 antigen on the MOLT-4V cell line. MOLT-4V cells were stained with a fluorescein-labelled 8F4 antibody (8F4-FITC) and investigated in flow cytometry (unfilled line, comparing with an isotype control (filled line)).

Fig. 12 shows the two-dimensional gel electrophoresis. A MOLT-4V cell lysate from  $300 \times 10^6$  cells was immunoprecipitated as described. The eluate was fractionated on a non-reducing SDS-PAGE (10% PAA), and the region around 60 kDa was cut out of the gel. To reduce the disulphide bridges in the 8F4 molecule, the piece of gel was incubated in 5.3 M urea, 0.5 M Tris, pH 8.0, 1% SDS, 1%  $\beta$ -mercaptoethanol at 50°C for 1 h, and the free cysteine residues in the molecule were alkylated with 10 mM iodoacetamide (Sigma, Deisenhofen) (37°C, 30 min). The piece of gel was equilibrated in 1xSDS-PAGE sample buffer for a further 30 min and mounted on a 12% PAA-SDS gel (with stacking gel). After fractionation by electrophoresis, the gel underwent silver staining. The location of the 8F4 protein was determined by surface iodination (cf. Fig. 1) and is marked by a circle. (All the procedures not described in detail were carried out by standard methods, see, for example, Westermeier, R., Electrophoresis in Practice, VCH Verlagsgesellschaft, Weinheim, 1997).

Fig. 13 shows a hybridization with Oligo 1 (SEQ ID NO:3). Lambda clones immobilized on nitrocellu-

lose filters were hybridized with Oligo 1 as described in the examples. Exposure on an X-ray film is depicted (detail).

Fig. 14 shows a Northern blot analysis with the 8F4 cDNA. Hybridization of a Northern blot with the 8F4 cDNA produces a band which migrates in the gel between the 18S and 28S RNA. Fig. 14A shows the behaviour as 2-signal-dependent (see above) activation antigen: no expression in resting lymphoid cells (PBL), strong expression in PMA+ionomycin-activated CD4+ T cells and distinctly reduced expression with PMA or ionomycin alone. Fig. 14B shows the strength of mRNA expression after different stimulation times (T cells (purified via nylon wool adherence, NTC), stimulated with PMA+ionomycin). Besides this the MOLT-4 cell lines (ATCC CRL-1582) which shows only minimal expression, and on the far right the MOLT-4V which was used for the cloning and which shows a distinct signal. Also loaded is the RNA from other cell lines on which no 8F4 expression was detectable in the analysis by flow cytometry: CEM (ATCC CCL-119), HUT-102 (ATCC TIB-162), HUT-78 (ATCC TIB-161), Jurkat (ATCC TIB-152), DG75 (Deutsche Sammlung von Mikroorganismen und Zellkulturen (DSMZ) ACC83), Karpas 299 (Fischer, P. et al. (1988), Blood, 72:234-240), DEL (Barbey, S. et al. (1990), Int. J. Cancer, 45:546-553).

Fig. 15 shows the amino acid sequence of the polypeptide 8F4 (SEQ ID NO:2).

Fig. 16 shows the 8F4 cDNA (SEQ ID NO:1).

The following examples illustrate the invention and are not to be understood restrictively.

#### Example 1: Generation of the 8F4 antibody

Balb/c mice were immunized with human T cells which had previously been activated for 24 h with 33 ng/ml of the phorbol ester phorbol myristate acetate (PMA) (Sigma, Deisenhofen) and with 200 ng/ml of the  $\text{Ca}^{2+}$  ionophore ionomycin (Sigma, Deisenhofen) (so-called "2-signal activation"). After boosting three times, the spleen cells of the mice were fused with the myeloma P3X63Ag8.653 (ATCC No. CRL-1580), and antibody-secreting hybridomas were generated by standard methods; cf. Peters and Baumgarten, Monoclonal Antibodies, Springer, Heidelberg, 1992. The resulting antibodies were screened for activated versus resting T cells

in flow cytometry. Activated ("2-signal activation") and resting T cells were incubated with the hybridoma supernatant and then labelled with a fluorescence-labelled secondary antibody; cf. Shapiro, Practical  
5 Flow Cytometry, Wiley-Liss, New York, 1995. Only the antibodies which recognize molecules which were induced exclusively by PMA and the  $\text{Ca}^{2+}$  ionophore ionomycin on the T-cell surface, but not by one of the agents alone ("2-signal molecules"), were selected for further puri-  
10 fication. The resulting antibodies were investigated in flow cytometry for similarity to or difference from known antibodies against activation molecules (cf. Table 1) on T cells. The criteria for this were, besides the abovementioned "2-signal dependence", the kinetics  
15 of induction on stimulated T cells and the expression on various cell lines.

#### Example 2: Immunoprecipitation of the 8F4 antigen

20 Surface molecules from activated human T cells were iodinated with  $^{125}\text{I}$  by standard methods and immunoprecipitated with the antibody 8F4 by standard methods; cf. Goding, Monoclonal Antibodies: Principle and Prac-  
tice, Academic Press, London, 1996. The antibody for  
25 the immunoprecipitation was coupled by the method of Schneider et al., *Journal of Biological Chemistry* 257 (1982), 10766-10769, to protein G (Pharmacia, Freiburg) (8F4 matrix). The matrix was washed as described by Schneider et al., see above. The immunoprecipitated 8F4  
30 molecule was analysed for its molecular mass in an SDS-PAGE (non-reduced and reduced) in a conventional way; Goding, see above.

#### Example 3: Flow cytometry

The 8F4-carrying T cells were analysed in flow  
35 cytometry by standard methods; cf. Shapiro, Practical Flow Cytometry, Wiley-Liss, New York, 1995.

Exemplary embodiment 3.1: Flow cytometry after induction of the 8F4 antigen on  $\text{CD4}^+$  T cells.

CD4<sup>+</sup> T cells from peripheral blood were stimulated with various agents in a conventional way, and investigated for expression of the 8F4 molecule in flow cytometry by a conventional method. The activation time for the T cells was between 24 hours and 144 hours with the various agents. Modes of activation: phorbol myristate acetate (PMA; 33 ng/ml), ionomycin (200 ng/ml), phytohaemagglutinin (PHA 1.5 mg/ml), OKT3 (monoclonal antibody against CD3), mixed lymphocyte reaction (MLR, between 50,000 CD4<sup>+</sup> T cells and 100,000 B cells), mAk 9.3 (monoclonal antibody against CD28), staphylococcal enterotoxin B (SEB, 0.1 ng/ml). Analysis revealed that various stimuli are suitable for inducing the 8F4 molecule on T cells, but the expression density differs. The most potent stimuli, besides the highly active pharmacological agents PMA and ionomycin, are those which represent a costimulatory situation such as, for example, accessory cells in the MLR or the costimulating mAk 9.3.

Exemplary embodiment 3.2: Kinetics of induction of the 8F4 antigen on CD4<sup>+</sup> T cells after activation with PMA and ionomycin.

CD4<sup>+</sup> T cells from peripheral blood were stimulated with PMA (33 ng/ml) and ionomycin (200 ng/ml) in a conventional way and investigated after 0, 4, 8, 12, 24 and 48 hours for expression of the 8F4 molecule by flow cytometry in a conventional way. The molecule is detectable on the surface after only four hours, and thus belongs to the class of relatively early activation antigens. There is still good expression of the antigen even after 48 hours.

Exemplary embodiment 3.3: Flow cytometry to identify molecules which are involved in the induction of 8F4 in the "mixed lymphocyte reaction".

50,000 CD4<sup>+</sup> T cells from peripheral blood were cocultivated with 100,000 allogeneic tonsillar B cells for 6 days (37°C, 5.2% CO<sub>2</sub>, 200 µl of RPMI 1640 with 10% FCS in 96-well round-bottom plates) and then investigated for expression of the 8F4 molecule in flow cytometry. At the start of cultivation, various antibodies (anti-CD80, anti-CD86, anti-MHCII; all 10 mg/ml) were added to the culture in order to examine the dependence of 8F4 induction on these molecules. Expression of 8F4 can be blocked only by blockade of the CD86/CD28 interaction, but not by blockade of CD80. The blockade effect in this case is even stronger than the blockade of MHCII (positive control).

Exemplary embodiment 3.4: Expression of 8F4 on T and B cells from human tonsils.

B cells and T cells from tonsillar tissue from various sources were purified in a conventional way and investigated by flow cytometry for expression of the 8F4 molecule. Whereas the signal was not unambiguously significant on B cells, there was expression of the 8F4 molecule in varying density by about 50-80% of tonsillar T cells. It is possible in this case to identify two populations differing in the level of fluorescence (8F4 high and low, respectively), and differing in expression on the various tonsils. Thus, for example, tonsils show a pronounced 8F4 low population and other tonsils show a pronounced 8F4 high population.

Exemplary embodiment 3.5: Coexpression of the 8F4 molecule with other activation markers.

T cells purified from human tonsils were analysed in 2-colour flow cytometry for coexpression of the 8F4 molecule with other activation markers. In tonsils, 8F4 is coexpressed with CD69 as well as with variants of the CD45 molecule. In this case, the 8F4 high cells are unambiguously correlated with a CD45R0 expression,

while the 8F4-negative cells carry the phenotype CD45RA. CD45RA is mainly expressed by so-called "naïve" T cells, whereas CD45RO is associated with an effector cell function. The 8F4<sup>+</sup> cells are thus mainly "mature" T cells. CD45RO and CD45RA are isoforms of CD45.

Example 4: Localization of 8F4-positive cells in the tonsil

Tonsillar tissue in frozen sections was stained with the 8F4 antibody in the APAAP technique (alkaline phosphatase-anti-alkaline phosphatase) by standard methods. 8F4<sup>+</sup> cells were found preferentially in the germinal centre of the tonsils, but also in part in the T-cell zone of the tonsils.

Example 5: Costimulation of T lymphocytes

96-well plates were coated with a goat anti-mouse Ig antibody (20 µg/ml), washed, and loaded with the anti-CD3 monoclonal antibody OKT3 (various dilutions of an ascites) and the 8F4 antibody according to the invention (2 µg/ml). The OKM1 antibody or the 2A11 antibody (both 2 µg/ml) were used as isotype control.

Exemplary embodiment 5.1: Enhanced expression of activation molecules on T lymphocytes after costimulation by 8F4.

Purified CD4<sup>+</sup> T cells from peripheral blood were activated with various concentrations of the monoclonal antibody OKT3 and, at the same time, costimulated with the 8F4 antibody or a nonspecific antibody of the same isotype. As comparison, costimulation was carried out with the anti-CD28 antibody-9.3, one of the strongest known costimulatory antibodies. Even with optimal stimulation by CD3, a costimulatory effect is still to be seen both with the mAb 8F4 and with the mAb 9.3. In the suboptimal OKT3 region, that is to say the region in which complete T-cell activation cannot be achieved

without costimulation, both antibodies are able to increase the expression of other activation antigens by a factor of 4 to 100, and the effect of the anti-CD28 antibody is still visible even at very high OKT3 dilutions. This is attributable to the fact that with very weak OKT3 stimulation the 8F4 antigen is no longer brought to the cell surface and thus cannot be crosslinked by the mAk 8F4 either.

Exemplary embodiment 5.2: Comparison of the costimulating effect of 8F4 with the costimulating effect of CD28.

Purified CD8<sup>+</sup> T cells were stimulated with a suboptimal concentration of the monoclonal antibody OKT3 for 51 h. The costimulators employed were antibody 8F4, antibody 9.3 (anti-CD28) and isotype controls (2 µg/ml each). After completion of the stimulation time, the T-cell proliferation rate was determined by <sup>3</sup>H-thymidine incorporation. In parallel cultures, the supernatant was removed and the concentration of the cytokines ATAC/lymphotactin and IL-2 was determined. 8F4 and CD28 differ greatly from one another in relation to IL-2 synthesis. CD28 costimulation leads, as also described in the prior art (Chambers and Allison, *Current Opinion in Immunology* 9 (1997), 396-404), to very extensive IL-2 secretion. By contrast, IL-2 production with 8F4 is below the detection limit. However, proliferation is comparable in the two mixtures, and thus the autocrine growth of the T cells must be attributed to other factors on costimulation of 8F4. The two antibodies also differ scarcely at all in the costimulatory effect in relation to secretion of the lymphokine ATAC.

Example 6: Determination of the immunoglobulins synthesized by B cells after interaction with 8F4-costimulated T cells



96-well plates were coated with a goat anti-mouse Ig antibody (20 µg/ml), and loaded with the anti-CD3 monoclonal antibody OKT 3 (1:500 to 1:80,000 ascites) and the 8F4 antibody according to the invention (2 µg/ml). The OKM1 antibody or the 2A11 antibody was used as isotype control. In some experiments, a costimulation was carried out with a CD28-specific antibody ("9.3") for comparison; cf. Hara et al., *Journal of Experimental Medicine* 161 (1985), 1513-1524. 50,000 purified (Magnetobeads, Dynal, Hamburg) CD4<sup>+</sup> T cells (>95% purity) from peripheral blood and 25,000 allogenic tonsillar B cells (negative selection by T-cell rosetting with sheep erythrocytes, 96% purity) were pipetted into each well of the culture plates pretreated in this way, and cocultivated for 8 days. After this period, the supernatant was removed and analysed for the concentration of secreted immunoglobulins of the IgM and IgG types in an ELISA in a conventional way; cf. Nishioka and Lipsky, *Journal of Immunology* 153 (1994), 1027-1036.

Exemplary embodiment 6.1: Enhancement of the synthesis of antibodies of the IgM and IgG types by the B cells after costimulation of T cells.

Purified CD4<sup>+</sup> T cells from peripheral blood were cocultivated with allogeneic B cells from tonsils for 8 days in a conventional way. With suboptimal stimulation of the T cells with the OKT3 antibody, the costimulation of the T cells by 8F4 enhances the secretion of IgM and IgG immunoglobulins by a factor of 40.

Example 7: Prevention of the activation-induced apoptosis of peripheral T cells after costimulation by 8F4.

Peripheral T cells (purified by nylon wool adherence in a conventional way), were stimulated with PHA (1.5 mg/ml) for 20 h and cultivated with IL-2 for 6 days. The cells were then restimulated by OKT3 with

and without costimulation by mAk 8F4 (2  $\mu$ g/ml). The apoptosis was determined by staining the DNA with propidium iodide in flow cytometry (FACS). With suboptimal stimulation via the T-cell receptor complex, costimulation by 8F4 can reduce the proportion of apoptotic cells by a factor of 4.

Example 8: Cloning of the cDNA coding for the 8F4 protein

A cell line (MOLT-4V) which expresses the 8F4 antigen constitutively was identified in flow cytometry by staining with a fluorescent dye-coupled 8F4 antibody (Fig. 11). The MOLT-4V line is a variant of the human T-cell line MOLT-4 (American Type Culture Collection (ATCC) CRL-1582).

This cell line was used for preparative purification of the 8F4 antigen with the aid of the monoclonal antibody:

The cells were cultivated on a large scale (150 l) in roller culture bottles and removed by centrifugation, and the cellular proteins were extracted using a lysis buffer (50 mM Tris, pH 8.0, 150 mM NaCl, 1 mM EDTA, 1 mM PMSF (Sigma, Deisenhofen), 1% NP-40 (Boehringer, Mannheim)). Cell nuclei and other insoluble constituents were removed by ultracentrifugation. The cell lysate obtained in this way was preincubated with Sepharose CL4-B (Pharmacia, Freiburg) for 2 h in order to remove proteins which bind nonspecifically to Sepharose. Incubation then took place with the 8F4 immunoaffinity matrix described in Example 2 above (4 h at 4°C). The matrix was packed into a column and then washed several times under conditions with which there is exclusive removal of nonspecifically binding proteins (1.50 mM Tris, pH 8.0, 300 mM NaCl, 1 mM EDTA, 1 mM PMSF, 0.5% NP-40; 2.50 mM Tris, pH 8.0, 150 mM NaCl, 1 mM EDTA, 1 mM PMSF, 0.5% NP-40, 0.1% SDS; 3.

0.2 M glycine pH 4.0, 0.5% CHAPS (Merck, Darmstadt)).  
The 8F4 antigen was eluted from the matrix with 0.2 M  
glycine, pH 2.5, 0.5% CHAPS. The eluate was concen-  
5     trated by ultrafiltration (Amicon Centricon 10, Milli-  
pore, Eschborn).

      In order to achieve further purification of the  
8F4 molecule, the dimeric structure of the molecule  
(see Fig. 1) was utilized in a two-dimensional gel  
electrophoresis (nonreducing/reducing): since most pro-  
10    teins occur as monomer, they migrate on a diagonal in  
gel electrophoresis, whereas the 8F4 molecule migrates  
at 55-60 kDa in the 1st dimension (nonreducing) and at  
27 and 29 kDa (Fig. 12) in the 2nd dimension (reduc-  
ing).

15       For preparative fractionation, the immunopre-  
cipitates from in each case  $20 \times 10^9$  cells were prepared  
as described above for Fig. 12 and fractionated in two-  
dimensional gel electrophoresis, the gel was stained  
with Coomassie blue G250 (Biorad, Munich) and the areas  
20    indicated in Fig. 12 were separately cut out of the gel  
(8F4-27 kDa and 8F4-29 kDa respectively).

      For peptide microsequencing, the proteins from  
in each case 4 pieces of gel were digested with trypsin  
and eluted from the gel. The tryptic fragments were  
25    fractionated by HPLC and individual fractions were sub-  
jected to Edman degradation (method described in detail  
in Groettrup, M. et al. (1996), Eur. J. Immunol.,  
26:863-869).

      Sequencing of the 8F4-29 kDa sample revealed, besides  
30    fragments of known proteins, a peptide sequence XRLTDVT  
for which no human correlate was found in any of the  
protein databases.

      Unambiguous translation back of a protein se-  
quence into a DNA sequence is not possible. Thus,

translation of the above peptide sequence back into an oligonucleotide with 17 nucleotides results in 2048 permutations. However, a specific method (Wozney, J.M. (1990), Methods Enzymol. 182:738-751) makes it possible to screen a cDNA bank with degenerate oligonucleotides. On the basis of the peptide sequence found, 2 oligonucleotides (Oligo 1 (SEQ ID NO:3); MGN CTS ACN GAY GTN AC, 512 permutations; Oligo 2 (SEQ ID NO:4): MGN YTD ACN GAY GTN AC, 1024 permutations) were synthesized.

For screening, a cDNA bank was constructed from the MOLT-4V cell line also used for the protein purification:

Complete RNA was isolated by the guanidinium/CsCl method (Chirgwin, J.M. et al. (1979), Biochemistry 18:5294-5299), and mRNA was concentrated on Oligo-dT-cellulose columns (Gibco BRL, Eggenstein). Synthesis of the first and second cDNA strands was carried out using a commercial cDNA synthesis system (Gibco BRL, Eggenstein) using Oligo-dT primers in accordance with the manufacturer's instructions. The cDNA was ligated via EcoRI adaptors into the Lambda ZAPII vector (Stratagene, Heidelberg).

The cDNA bank was plated out by standard methods (Vogeli, G. and Kaytes, P.S. (1987), Methods Enzymol., 152:407-515) and the Lambda DNA was immobilized on nitrocellulose filters (Optitran BA-S 85, Schleicher & Schuell, Dassel).

The abovementioned oligonucleotides were radio-labelled using T4 polynucleotide kinase (NEBL, Schwalbach) and  $\gamma$ -<sup>32</sup>P ATP (NEN Du Pont, Brussels) (Wallace, R.B. and Miyada, C.G. (1987), Methods Enzymol., 152:432-442).

Hybridization of the filters took place in a buffer described for degenerate oligonucleotides (Wozney, J.M. (1990), Methods Enzymol. 182:738-751) with 3 M tetramethylammonium chloride (Roth, Karlsruhe) at 48°C. The filters were washed as described in the abovementioned reference, the washing temperature being 50°C. Exposure of these filters on an X-ray film revealed about 50 positive clones per 100,000 plated phages (Fig. 13).

6 clones were further characterized by transferring them by in vivo excision, using the method described by the manufacturer of the vector (Stratagene, Heidelberg), into a plasmid vector, and partially sequencing with T3 and T7 primers (BigDye Terminator Cycle Sequencing Kit, Applied Biosystems, Foster City, USA). One of the clones contained a sequence which on translation provided exactly the peptide sequence which was sought. This clone was used for hybridization of a Northern blot (Fig. 14) (Kroczeck, R.A. (1993), J. Chromatogr., 618, 133-145). The expression pattern of the mRNA corresponded exactly to the expression of the 8F4 molecule as was known from investigations on the monoclonal antibody by flow cytometry. Since the clone which was found contained only the 3' end of the cDNA sought, a fragment on the 5' side was used to isolate the complete 8F4 cDNA. Several clones were sequenced on both strands.

The 8F4 cDNA (2641 nucleotides) is depicted in Fig. 16 and in the sequence listing under SEQ ID NO:1, and codes for a protein having 199 amino acids (Nucleotides 68-664), depicted in Fig. 15 and in the sequence listing under SEQ ID NO:2. Sequencing of several independent clones from the cDNA bank showed some deviations from the sequence shown here, but these are all in the 3'-untranslated region:

Pos. 909-910:deletion

Pos. 1631:T->C

Pos. 2074:G->T

Pos. 2440:G->C

5 Pos. 2633: alternative polyadenylation site

111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158  
2159  
2160  
2161  
2162  
2163  
2164  
2165  
2166  
2167  
2168  
2169  
2170  
2171  
2172  
2173  
2174  
2175  
2176  
2177  
2178  
2179  
2180  
2181  
2182  
2183  
2184  
2185  
2186  
2187  
2188  
2189  
2190  
2191  
2192  
2193  
2194  
2195  
2196  
2197  
2198  
2199  
2200  
2201  
2202  
2203  
2204  
2205  
2206  
2207  
2208  
2209  
2210  
2211  
2212  
2213  
2214  
2215  
2216  
2217  
2218  
2219  
2220  
2221  
2222  
2223  
2224  
2225  
2226  
2227  
2228  
2229  
2230  
2231  
2232  
2233  
2234  
2235  
2236  
2237  
2238  
2239  
2240  
2241  
2242  
2243  
2244  
2245  
2246  
2247  
2248  
2249  
2250  
2251  
2252  
2253  
2254  
2255  
2256  
2257  
2258  
2259  
2260  
2261  
2262  
2263  
2264  
2265  
2266  
2267  
2268  
2269  
2270  
2271  
2272  
2273  
2

Table 1:

Table 1 summarizes the antibodies used (clone), their source of origin (source), the specificity for their particular antigen (specificity) and, where appropriate, their labelling (label).

Specificity	Label	Isotype	Clone	Source
CD3	Cy-Chrome	IgG1	UCHT1	Pharmingen, Hamburg
CD3	-	IgG2a	OKT3	ATCC CRL-8001
CD11b	-	IgG2b	OKM1	ATCC CRL-8026
CD25	FITC	IgG2a	B1.49.9	Immunotech, Hamburg
CD28	-	IgG2a	9.3	Immunex Corp., Seattle
CD45RA	Cy-Chrome	IgG2b	HI100	Pharmingen, Hamburg
CD45RO	FITC	IgG2a	UCHL1	Immunotech, Hamburg
CD69	FITC	IgG1	FN50	Pharmingen, Hamburg
CD80	-	IgG1	L307.4	Becton Dickinson, Heidelberg
CD86	-	IgG2b	IT2.2	Pharmingen, Hamburg
CD154	FITC	IgG1	TRAP-1	Hybridoma <sup>1</sup>
MHCII	-	IgG2a	L243	ATCC HB-55
8F4	-	IgG1	8F4	Hybridoma <sup>1</sup>
8F4	Biotin	IgG1	8F4	Hybridoma <sup>1</sup>
Isotype IgG1	-	IgG1	2A11	Hybridoma <sup>1,2</sup>
Isotype IgG1	FITC	IgG1	2A11	Hybridoma <sup>1,2</sup>
Isotype IgG1	Biotin	IgG1	ASA-1	Hybridoma <sup>1</sup>

<sup>1</sup> The hybridoma cell line was generated in a conventional way, and the antibody was purified and labelled where appropriate.

<sup>2</sup> Directed against a synthetic peptide

5

The antisera and secondary reagents used in the examples were purchased from: goat anti-mouse Ig, FITC conjugated, from Jackson Immuno Research Lab., USA; Streptavidin, PE-conjugated, from Jackson Immuno Research Lab., USA; rabbit anti-mouse Ig fraction, from Sigma, Deisenhofen.

15 It will be apparent to those skilled in the art that various modifications and variations can be made to the compositions and processes of this invention. Thus, it is intended that the present invention cover such modifications and variations, provided they come within the scope of the appended claims and their equivalents.

20 Priority applications DE 19741929.1 filed September 23, 1997 and DE 19821060.4 filed May 11, 1998 including the specification, drawings, claims and abstract, are hereby incorporated by reference. All publications cited herein are incorporated in their entireties by  
25 reference.